



nutrace




Skretting Australia

Residue Monitoring

Summary of Results 2009-2013
Issued April 2014



About Skretting



Skretting is a world leader in nutritional solutions for farmed fish and shrimp.

Total annual production and supply of high quality feeds is approximately 1.7 million tonnes (2012). Skretting has operating companies on five continents to produce and deliver feeds from hatching to harvest for more than 60 species of farmed fish and shrimp. Skretting is owned by international animal nutrition and fish feed group Nutreco.

Nutreco is a global leader in animal nutrition and fish and shrimp feed.

Experience across 100 years brings Nutreco a rich heritage of knowledge. Nutreco employs approximately 10,000 people in 30 countries, with sales in 80 countries. Nutreco is listed on the NYSE Euronext stock exchange in Amsterdam.

Read more at www.nutreco.com

Quick Links



About residue testing

Skretting conduct regular testing of our feeds for undesirable substances.

This is part of Skretting's Quality System, which acts to validate the quality controls performed throughout the year. These controls include frequent analysis of raw materials, supplier assessments and systems to control the pellet manufacturing process.

Skretting's Food Safety Team regularly review potential residues based on a risk assessment considering the scale of use, toxicity and persistence of each compound.

A global testing program for these residues is set annually. Testing is conducted by Skretting-approved, accredited laboratories that have demonstrated the highest level of competency and repeatability.

Skretting Australia undertakes residue testing on both raw materials and finished feeds to ensure their quality.

Results reported here relate to complete fish feeds produced by Skretting Australia. Samples chosen best represent production since publishing the previous Residue Monitoring Report in 2013.

“Skretting Australia undertakes residue testing on both raw materials and finished feeds to ensure their quality”

Quick Links



About this report

This Residue Monitoring Report summarises the level of undesirable substances in Skretting Australia feeds from 2009 to 2013.

Results from monitoring in 2013 indicate that Skretting Australia feeds have again met all Australian and European requirements and that the levels of undesirable substances found in feeds are substantially less than the limits set by authorities.

Skretting's Residue Monitoring Report uses the Australian residue limits (maximum residue limit and

the extraneous residue limit MRL/ ERL) as set by the **Australian Pesticides and Veterinary Medicines Authority (APVMA)**, as well as the EU statutory limits (equivalent to MRL standards) where they are available. If an Australian residue limit does not exist for a parameter, the relevant EU statutory limit has been used.

We provide this report to keep Skretting customers informed of the status of our monitoring results.

To view previous reports, [click here](#)

Results from monitoring in 2013 indicate that Skretting Australia feeds have again met all Australian and European requirements

Quick Links



Feed to Food Quality & Safety

Nutrace® is Skretting's unique, global feed-to-food quality and safety system. It ensures consistency throughout the production process, from raw materials to final feed solutions.

Built on five strong pillars, Nutrace® ensures that our customers and end consumers can have full confidence with regard to feeds in the farmed fish value chain.

In 2013 Skretting Australia became the 3rd Skretting company to be awarded Nutrace® compliance.

[Click here to read more about Nutrace®](#)



Certified Quality
& Food Safety



Ingredient and Supplier
Assessment & Management



Monitoring
& Control



Risk Management



Tracking & Tracing

Quick Links



PCBs & Dioxins

BACKGROUND

Polychlorinated biphenyls (PCBs) are extremely persistent organic pollutants historically used as coolants, plasticisers, lubricants etc. The term “dioxins” includes polychlorinated dibenzo-p-dioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs). These are listed on laboratory reports as PCDD/F and are the main focus in terms of food safety. PCDD/Fs are unwanted by-products of chemical manufacture, bleaching processes and all combustion processes.

HEALTH ISSUES

PCBs and dioxins are toxic molecules, which in small quantities may cause problems with reproduction and cancer. PCB exposures, particularly before birth, have been linked in humans with lower IQ, hyperactivity, shortened attention span and delayed acquisition of reading skills.

LIMITS

Substance	Unit	Australia ¹	EU/ Norway ²
Dioxins (Dioxins & furans)	TEQ (WHO) ng/kg	No	1.75
Sum of Dioxin & Dioxin-like PCBs	TEQ (WHO) ng/kg	No	5.5

ng = nanogram

kg = kilogram

TEQ = Toxic Equivalent (the amount of toxin or other poison per kilogram of body weight necessary to kill an animal)

WHO = World Health Organisation (Standard)

1. The MRL Standard: Maximum residue limits in food and animal feedstuff. APVMA December, 2012 32pp
2. OJ L 91, 29.3.2012, p.18-20. Commission Regulation (EU) No 277/2012

Quick Links




Polychlorinated biphenyls (PCBs) are released into the environment and eventually enter the food chain and can build up in fish and other marine animals. Manufacturing of PCBs has been banned in most countries since the 1980s. Australia banned the importation of PCBs in 1975.

Dioxins have been shown to bioaccumulate in humans and animals due to their lipophilic (fat-liking) properties. Dioxins enter the general population almost exclusively from ingestion of food, specifically through the consumption of meat and dairy products.


PCBs and dioxins (and furans) have a similar structure and they therefore have a similar toxic effect. The PCBs are usually present in much higher quantities than dioxins, but are less toxic. The most toxic PCBs are classed as 'dioxin-like' by the **World Health Organisation (WHO)**. It is these 12 'dioxin-like' PCBs which the EU has set limits for in combination with dioxin residues.

In the figures to the right, "ng/kg" denotes the toxic equivalent (WHO) including limit of quantitation (LOQ) fat weight ng/kg.

PCDD & PCDF



SUM DIOXINS & DIOXIN-LIKE PCBs



Pesticides

BACKGROUND

Pesticides have been widely and commonly used to protect crops, livestock, buildings and households from pests. Although widespread use has been banned in many countries, several pesticides are still produced in developing countries. Many pesticides continue to be detected in precipitation, soil, sediment, biota, aquatic ecosystems and food.

HEALTH ISSUES

The health effects of pesticides depend on the type of pesticide. Some, such as the organophosphates and carbamates, affect the nervous system. Others may irritate the skin or eyes. Some pesticides may be carcinogenic. Others may affect the hormone or endocrine system in the body.

LIMITS

Pesticide	Unit	Australia ¹	EU/ Norway ²
Aldrin & Dieldrin (sum of HHDN & HEOD)	mg/kg	E 0.01	0.02
Chlordane (sum of cis- trans- & oxy-chordane isomers)	mg/kg	E 0.01	0.02
DDT (sum of o,p'-DDT; p,p'-DDE; p,p'-DDT & p,p'TDE)	mg/kg	E 0.05	0.05
Endosulfan (sum of alpha- & beta-endosulfan & endosulfan sulphate)	mg/kg	0.3	0.005
Endrin (sum of endrin & delta-keto endrin)	mg/kg	E 0.03	0.01
Heptachlor (sum of heptachlor & heptachlor epoxide)	mg/kg	E 0.02	0.01
Hexachlorobenzene (HCB)	mg/kg	E 0.01	0.01
alpha-Hexachlorocyclohexane (alpha-HCH)	mg/kg	No	0.02
beta-Hexachlorocyclohexane (beta-HCH)	mg/kg	No	0.01

mg = milligram

kg = kilogram

1. The MRL Standard: Maximum residue limits in food and animal feedstuff. APVMA December, 2012 32pp. 'E' denotes an Extraneous Residue Limit (ERL)

2. OJ L140, 30.5.2002, p. 10. Directive 2002/32/EC of the European Parliament and of the Council of 7 May 2002 on undesirable substances in animal feed

In previous years, samples were screened for total pesticides, but from 2013 onwards each pesticide is analysed individually. The Residue Report will now show actual level, not just whether the sample was under the threshold.

Quick Links




Aldrin and dieldrin (a metabolite of aldrin as well as a marketed pesticide) are both fat soluble, persistent and bio-accumulating organochlorine insecticides.


Widely used as insecticides in agriculture, the registration of the last aldrin and dieldrin products in Australia were cancelled in 1994 and 1988 respectively. Both compounds are classified as Schedule 6 ('Poison') in the [Standard for the Uniform Scheduling of Medicines and Poisons \(SUSMP\)](#).

Worldwide, the use of both compounds is severely restricted or banned in many countries. In the environment, aldrin is rapidly converted to dieldrin.

ALDRIN



DIELDRIN




*Not detected LOQ <0.002

Chlordane is a non-systemic (ie not taken up in the plant) insecticide of agricultural crops. It is a restricted chemical product and is classified as Schedule 6 (“Poison”) in the SUSDP.

Chlordane is not used in Australia - the last registered Chlordane product was withdrawn in 1997. Chlordane is banned in Europe (1981) and the USA (1978) and currently in most other countries worldwide.

Chlordane is highly insoluble in water, persistent, bio-accumulating and highly lipophilic. In accordance with the Australian residue limit (ERL) standard, Chlordane is reported as the the sum of cis-, trans- and oxy-Chlordane isomers.

CHLORDANE




Dichlorodiphenyltrichloroethane (DDT) is an insecticide that was widely applied in agriculture and forestry. DDT products were classified as Schedule 5 (‘Caution’) or Schedule 6 (‘Poison’) in the SUSDP.

Since the early 1970s severe restrictions and bans on its use have been introduced in many countries (eg. Banned in the USA in 1972 and Europe since 1986). In Australia, registrations of all DDT products had been cancelled by the late 1990’s (the majority of products had not been used since the mid-1980’s).

DDT is highly insoluble in water, lipophilic and persistent in the environment. Because of the lipophilic properties and persistence in the environment, DDT and related compounds are bio-accumulating and biomagnified along the food chain. Although it is banned in most countries worldwide, DDT is still used for vector control (eg mosquitoes), especially in areas with endemic malaria.

DDT



*Not detected LOQ <0.002

Endosulfan is a non-systemic organochlorine pesticide used in agricultural and horticultural crops for control of insects and mites.

Registration of endosulfan in Australia was cancelled in October 2010. Endosulfan is banned in Europe (2006) and currently restricted or banned in most other countries worldwide.

In contrast to the majority of organochlorine pesticides, endosulfan is less lipophilic – consequently, biomagnification and bio-accumulation along the food chain is less likely to occur.


In accordance with the Australian residue limit (MRL) standard, endosulfan is reported as the sum of alpha- & beta-endosulfan isomers and endosulfan sulphate.

Endrin is a fat soluble organochlorine insecticide - at one time registered in Australia as an insecticide, miticide and avianicide.


Classified as Schedule 7 ('Dangerous Poison') in the SUSDP, the last Australian-registered endrin product was cancelled in 1990. Endrin has been banned in most countries worldwide during the last 25 years.

Endrin is partly transformed in the environment into delta-keto endrin.

ENDOSULFAN



ENDRIN



*Not detected LOQ <0.002

Heptachlor is a non-systemic contact insecticide – at one time registered in Australia as a termatocide and insecticide. Classified as Schedule 6 ('Poison') in the SUSDP, the last Australian-registered heptachlor product was cancelled in 1997 (the majority of heptachlor products were cancelled by the end of 1990). Heptachlor is banned in Europe (1984) and most other countries worldwide.


In the environment, heptachlor breaks down to heptachlor epoxide and photoheptachlor. All these compounds are lipophilic, persistent and bio-accumulate in the food chain.

In accordance with the Australian residue limit (ERL) standard, heptachlor is reported as the the sum of heptachlor and heptachlor epoxide.


Heptachlorobenzene (HCB) is an agricultural pesticide used as a fungicide (seed disinfectant). The last Australian-registered HCB product was banned in 1987. HCB is banned in Europe (1981) and most other countries worldwide.

HCB is highly insoluble in water, lipophilic, persistent and bio-accumulates in the food chain.

HEPTACHLOR



HEXACHLOROBENZENE (HCB)




*Not detected LOQ <0.002


Technical Hexachlorocyclohexane (HCH) was used as an insecticide worldwide. It is a mixture of isomers – the four predominant are alpha-, beta-, delta- and gamma-HCH (also known as lindane).

Technical HCH is banned in Europe (1978) and most other countries worldwide. Alpha- & beta-HCH are lipophilic, persistent in the environment, bio-accumulating and biomagnified along the food chain.

ALPHA-HCH



BETA-HCH



*Not detected LOQ <0.002

Heavy Metals

BACKGROUND

Heavy metal pollution commonly arises from the purification of metals and unlike organic pollutants, heavy metals do not decay. Emissions of heavy metals such as mercury, lead, cadmium and arsenic into the environment occur via a wide range of processes and pathways including the air, surface water, and soil.

HEALTH ISSUES

One of the largest problems with the persistence of heavy metals in the environment is the potential for bioaccumulation and biomagnification causing heavier exposure for some organisms than is present in the environment alone. If the human body is exposed to heavy metals of a specific concentration for an interval of time then this can cause serious illness.

LIMITS

Substance	Unit	Australia ¹	EU/ Norway ²
Arsenic	mg/kg	No	10.0
Cadmium	mg/kg	No	1.0
Lead	mg/kg	No	5.0
Mercury	mg/kg	No	0.2

mg = milligram

kg = kilogram

1. The MRL Standard: Maximum residue limits in food and animal feedstuff. APVMA December, 2012 32pp

2. OJ L140, 30.5.2002, p. 10. Directive 2002/32/EC of the European Parliament and of the Council of 7 May 2002 on undesirable substances in animal feed

Quick Links




Arsenic and its compounds are used as pesticides and in various alloys. The toxicity of arsenic is strongly dependent on its chemical form. Although inorganic arsenic is highly toxic, organic arsenic is not.

Accumulated arsenic in fish and shellfish is predominantly in organic form, and aquatic animals show a wide range of sensitivities to different arsenic compounds.


Arsenic in fish feed is also predominately in the organic form, which is much less toxic than the inorganic form.

Cadmium is commonly found in its metallic form and as sulfides and sulfates. Globally, about three-quarters of cadmium is used in batteries and most of the remaining quarter is used mainly for pigments, coatings and plating, and as stabilizers for plastics.

ARSENIC




CADMIUM




Sources of lead found in the environment are multiple, and the metal is truly ubiquitous, being commonly found in food, water, and air. Evidence exists that lead in the environment has increased during the past 200 years, and it is not surprising that it can be found as a contaminant of aquatic animals.


LEAD



Mercury is much more harmful to living organisms as an organic metal compound than as the element. The most toxic form of mercury is methylmercury, which damages the central nervous system. It is found in fish because industrial effluents containing mercury are discharged into rivers or seas where the mercury is converted into methylmercury by bacteria. It then moves up the food chain and accumulates in the bodies of some large wild fish, such as shark, marlin and swordfish. In farmed salmon the levels of mercury are very low, almost not detectable.

MERCURY





www.skretting.com.au